Presentation 2 - Kathleen Considine and Vince Iannacchione



RTI Areas of Expertise Survey Methodology CAI Systems GIS Experts Database Management Project Management Project Management Psychometrics Database Management Sio-Specimen Collection Statistical Analysis

Primary Analytic Objectives

Among Gulf War Era Veterans

- Estimate the prevalence of:
 - any Haley syndrome
 - each of the three primary Haley syndromes
- Test for differences between those who were:
 - Deployed to Kuwaiti Theater of Operations (KTO)
 - Not deployed to KTO (but deployable)

www.rti.org

Secondary Analytic Objectives

- Validate and refine the factor analysis model used to derive original Haley syndromes
- Estimate magnitude of the "Healthy Warrior Effect"

"Bias caused by the disproportionate concentration of prewar illness in the non-deployed population."

Study Design Overview

Three-Phase Study of Gulf War Era (1990-91) Veterans

Phase 1: Approximately 8,000 Telephone Interviews (CATI)

Phase 2: Approximately 2,000 Blood Specimens

Phase 3: Approximately 400 Clinic Exams

www.rti.org

Components of the Interview Phase

- CATI Questionnaire Development
- CATI Programming
- Telephone Interviewing Protocol
- Quality Assurance

Phase 1 – CATI Questionnaire

Questionnaire includes three modules

- Symptoms (771 questions)
- Exposures (244 questions)
- Family Issues (76 questions)

Average length of interview

- 60 minutes for those healthy and not deployed to KTO
- 2.5 hours for those deployed with health issues

www.rti.org

CATI Questionnaire

Symptoms Module contains questions regarding:

- Case definitions of Haley, Steele, Fukuda, Kang, CFS (CDC), FM (survey) and MCS (Iris Bell's Chemical Odor Intolerance scale).
- Physical and mental functional status (SF-12)
- Physician-diagnosed conditions
- Medical assessments of vision and balance disturbance
- Postwar driving abilities and record
- Health problems before the war
- Treatments tried and assessments of their effectiveness
- Time course of post-war symptoms

CATI Questionnaire

Exposures Module contains questions regarding:

- Chemical weapons alarms, Khamisiyah
- Use of insect repellents, pesticide-treated uniforms, flea collars
- Took pyridostigmine bromide (PB) and resulting side effects
- Vaccinations, medicines taken before and during deployment
- Exposure to oil fire smoke, jet fuel, other petroleum products
- Exposure to depleted uranium (CHPPM categories)
- Combat exposure scales (Keane; Haley)
- Individual location/date information if separated from unit
- Sexual harassment/assault

www.rti.org

CATI Questionnaire

Family Issues Module contains questions regarding:

- Fertility
- Outcome of each pregnancy
- Birth defects
- Evidence of learning disorders in children

10

CATI Questionnaire

Methodological Issues in Questionnaire Development

- Focus groups
- Adapt from paper-and-pencil (PAPI) to CATI
- Revisions based on other validated questions
- Methods for asking sensitive questions
- Techniques to minimize recall bias
- Identify programming and question wording issues
- Timing Test
- Procedures to alleviate mental fatigue

11

www.rti.org


CATI Programming

During the interview, the CATI program must be able to

- Calculate Haley, Steele, and Fukuda syndrome scores
- Match a veteran's unit with the Troop Location Data Base
- Schedule a continuation call
- Select a random sub-sample of non-syndromic veterans for blood specimens

12





Locating Veterans Initial batch tracing for all sample members Interactive tracing for unlocatable sample members Training CATI Interviewers 4 day-long interviewer trainings Interviewer training on dealing with refusals Interviewers must pass 2 certifications prior to beginning work



Endorsements and Study Clearances


Endorsements

Veterans Service Organizations

Clearances

- IRB Approval from RTI and UT-Southwestern
- HSRRB (DoD) clearance from Ft Detrick
- Certificate of Confidentiality from NIEHS

16





Quality Assurance Questionnaire Real-time silent monitoring Quality circle meetings with interviewers Interviewer Retraining Re-emphasize original training, address new issues Pilot Study Tested all facets of the design including data analysis

Phase 2 – Blood Specimens

Purpose

Determine association between Haley syndromes and PON level

Protocol

- Phase 2 participants selected from CATI respondents
 - All identified as syndromic using Haley definition or from groups of special interest
 - A random sub-sample from others
- Contract with EMSI to collect samples
- Samples sent to UT-Southwestern for analysis

20

www.rti.org



ORTI

Phase 3 – Clinic Exam

Purpose

Clinical verification of Haley syndromes

Protocol

- Phase 3 selected after Phase 2 completed from respondents
 - Case/control sample balanced on characteristics
- RTI will recontact Veterans by phone to gain cooperation
- UTSW will finalize appointments/transportation
- Clinic exam will take place at UTSW and will last 2 full days

21

www.rti.or



Time Line

Submitted HSRRB clearance for Pilot: Nov, 2003

Received HSRRB clearance for Pilot: Oct, 2004

Conducted CATI phase of the Pilot Survey: Jan, 2005 - Apr, 2006

Conduct blood phase of the Pilot: Nov – Dec, 2006

Obtain clearances for Main Study: Dec, 2006

Conduct Main Study: Jan – Dec, 2007

22

Survey Population

Target Population

3.75 million Gulf War Era veterans on active duty or in the reserves

- 752K veterans who were deployed to KTO
- 3M non-deployed veterans

Sampling Frame

- Created by Defense Manpower Data Center (DMDC)
- Updated KTO deployment status from CHPPM

23

www.rti.org

Stratified Sampling Design

Strata for CATI Phase

Deployed to KTO Location in KTO

Age Group Gender

Race Active or Reserve status

Special Strata

Strata for Blood Specimen Phase

Syndromic/Non-Syndromic

24

Stratified Sampling Design

Special Strata

- 24th Reserve Naval Construction Battalion
- Parents of children with Goldenhar Complex
- Twin Pairs
- Air crew members
- Aircraft maintenance personnel

25

www.rti.org

Design Optimization

Objective

Detect true differences between major reporting domains with acceptable statistical power.

Procedure

We will use the Sample Planning Tool developed by RTI staff for cost/variance optimization.

26

Sampling Weights

Sampling weights needed to adjust sample estimates for

- Selection bias
- Nonresponse bias

Weights calculated in a two-step process

- 1. Initial weight = inverse of selection probability
- 2. Nonresponse adjustment applied to initial weight

27

www.rti.org

Preliminary Sample Sizes

Selected: 13,000

Located: 11,400 (88%)*

Eligible: 11,000 (97%)*

CATI Participants: 8,000 (73%)*

* Pilot study results

For methodological use only. Estimates are not statistically significant.


28


Recommendations to Improve Response Rates


For the main survey, we will


- Use IRS information to help locate veterans
- Remove 'Gulf War' from the study title
- Address legal concerns about participating
- Maintain a web site for the study
- Promote survey in VSO publications
- Welcome other ideas!


_











Healthy Warrior Effect

Two Reasons for Non-Deployability

- **1. Medical**: An illness or physical condition (other than pregnancy) that prevented deployment to KTO.
- 2. Job Related: Military job not usually deployable or unit not fully prepared for deployment.

Pilot Study Respondents Not Deployed to KTO

Deployable: 19 (33%)

Non-Deployable:

Medical 5 (9%) Job Related 34 (58%)

For methodological use only. Estimates are not statistically significant.

35

www.rti.org



Pilot Respondents with Any Haley Syndrome

	Any Haley Syndrome		
Total #	#	% (wgt'd)	
60	13	17%	
19	2	5%	
34	1	5%	
5	2	15%	
	60 19 34	Total # # 60 13 19 2 34 1	

For methodological use only. Estimates are not statistically significant.

BRTI

Deployment and Any Haley Syndrome					
	75		,		
	Deployment status		Any Haley Syn*		
	Deployed	Nondep	Yes	No	
Total subjects	60	58	13	47	
Fibromyalgia (ACR screening def.)	10%	3%	46%	0%	
Chemical Odor Intolerance (Bell)	10%	3%	23%	6%	
MCS Screener (ATSDR)	27%	12%	54%	19%	
Severe MCS	3%	0%	8%	2%	
Chronic Fatigue Syndrome	1%	0%	0%	2%	
			*In depl	oyed only	

